

# Coastal NeuroSurgery<sup>P.A.</sup>

RICHARD C. HARTWELL, M.D., Ph.D.  
*Diplomate American Board of Neurological Surgery*

[www.CoastalNeuroSurgeryNJ.com](http://www.CoastalNeuroSurgeryNJ.com)

9 Hospital Drive Suite C10  
Toms River, New Jersey 08755  
Telephone (732) 341-1881  
Facsimile (732) 505-4453

## POSTERIOR CERVICAL DECOMPRESSION WITH FUSION CONSENT FORM

Name:

You have been scheduled for a posterior cervical decompression with fusion. Your surgery is scheduled for \_\_\_\_\_. The surgery involves making an incision in the back of your neck and removing the bone and ligament over the spinal canal and nerves in order to free up the spinal cord and nerves that are being compressed in your spinal canal. The surgery may also involve removing part of one or more disks if it is found that it is pressing on your nerves. Once your spinal cord and nerves are free, the fusion portion of the surgery will take place. Bone grafts, which are obtained from the bones removed from your spine, will be placed into the joint spaces of the unstable bones and screws will be placed into the bones and rods will be attached to the screws to hold the bones in place. The screws and rods are permanently placed and typically are not removed except in the case of infection or hardware breakage. You will wear a hard cervical collar for a minimum of 6 weeks. If narcotic pain medications are necessary I will prescribe them only in the immediate pre-operative treatment time leading up to the surgery and in the immediate post-operative recovery period but I will not continue them for more than 3 months following the surgery.

As with any surgery, there are risks that may occur during the surgery and in the postoperative period, including but are not limited to:

1. Blood loss and the need for transfusion: If you would like, you may donate your own blood to the Red Cross prior to your surgery. If you require a transfusion and did not donate any blood, the donated blood is carefully screened for AIDS (HIV) and hepatitis, but there are risks of you developing such infections from a transfusion.
2. Infection: There is a risk for infection. Antibiotics will be given to you right before the surgery and for at least 24 hours postoperatively in order to minimize the risks for infection.
3. There are risks for paralysis, nerve injury, loss of bowel, bladder, or sexual function, which may be temporary or permanent.
4. Persistent symptoms, worsening of symptoms or lack of benefit from the surgery: As with any surgery there is never a 100% guarantee that all or any of your symptoms will be completely resolved. There may already be permanent damage to your nerves, which may not improve at all in the postoperative period.
5. Cerebral spinal fluid leaks: During the surgery the covering over the nerves (dura) may tear and cause a leakage of spinal fluid. Typically, the tear is repaired with a suture during the surgery, however, the leak may persist after the surgery or a leak may occur which was not identified during the surgery. This situation may cause headaches, drainage of spinal fluid from the incision and possibly meningitis should the fluid become infected. Treatment for postoperative spinal fluid leakage includes laying flat in bed, IV fluids and possible placement of a spinal drainage catheter. It is rare that another operation is required to find the source of the leak.
6. Loss of mobility of the fused portion of your spine causing instability of the spinal segments above and below the level of your fusion in the future, which may require further surgery.

7. Infection/pain at the pinhole sites: During the surgery your head will be immobilized in a holder, which is secured to your skull with pins. There is a possibility that the pinhole sites can become infected or that you may have pain at these sites for a period of time after your surgery.
8. Misplacement of screws, breakage of screws or rods requiring re-operation in the future if symptoms occur.
9. Failure of a bony fusion to occur (nonunion/pseudoarthrosis) requiring re-operation.
10. Deep vein thrombosis (blood clot in the legs), pulmonary embolism.
11. Formation of a blood clot over the spinal cord (epidural hematoma) requiring re-operation.
12. Pressure sores over the chest wall, breasts, nipples and iliac crest with skin blisters and burns that may occur from the positioning on the OR table even though every effort is made to pad these areas. These may be temporary or permanent.
13. Pressure injury to nerves in the brachial plexus (brachial plexitis) and lateral femoral cutaneous nerves (meralgia paresthetica), which may occur from position on the OR table even though every effort is made to pad these areas. This may be temporary or permanent.
14. Risks of anesthesia: Adverse reaction to anesthesia given or any medication given during the surgery.
15. Heart attack, stroke, coma and death.

I acknowledge that I have read the above consent form and all options and alternative treatments were discussed with me by Dr. Hartwell. In addition, all of the above risks were discussed with me in detail, in laymen's terms, by Dr. Hartwell and I understand all the above risks and possible complications and wish to proceed with surgery.

1. For one week prior to the surgery: no Aspirin, Plavix, clopidogrel, Coumadin, warfarin, Fish Oil, Flax seeds, nutrient supplement pills, Vitamin E, Co-Q-10, Lovaza or anti-inflammatory medications such as Advil, Motrin, Aleve, Ibuprofen or Naprosyn one week prior to surgery.
2. No medicines for erectile dysfunction (ED medicines) 48 hours prior to surgery.
3. Nothing to eat or drink after 12:01 a.m. on \_\_\_\_\_.
4. Take the following medicines on the day of surgery with a small sip of water.

Signed: \_\_\_\_\_

Date: \_\_\_\_\_